

Phase Transitions in the Two-Dimensional Random Gauge XY Model

Petter Holme*

Department of Physics, Umeå University, 901 87 Umeå, Sweden

Beom Jun Kim†

Department of Molecular Science and Technology, Ajou University, Suwon 442-749, Korea

Petter Minnhagen‡

*NORDITA, Blegdamsvej 17, DK-2100, Copenhagen, Denmark and
Department of Physics, Umeå University, 901 87 Umeå, Sweden*

The two-dimensional random gauge XY model, where the quenched random variables are magnetic bond angles uniformly distributed within $[-r\pi, r\pi]$ ($0 \leq r \leq 1$), is studied via Monte Carlo simulations. We investigate the phase diagram in the plane of the temperature T and the disorder strength r , and infer, in contrast to a prevailing conclusion in many earlier studies, that the system is superconducting at any disorder strength r for sufficiently low T . It is also argued that the superconducting to normal transition has different nature at weak disorder and strong disorder: termed Kosterlitz-Thouless (KT) type and non-KT type, respectively. The results are compared to earlier works.

PACS numbers: 64.60.Cn, 64.70.Pf, 74.60.Ge, 74.76.Bz

I. INTRODUCTION

The XY gauge glass model¹ has attracted much interest in connection to the vortex glass phase of high- T_c superconductors². In three dimensions (3D) there is a general consensus that the XY gauge glass model exhibits a finite-temperature glass transition^{1,3,4,5}. However, in 2D there exist conflicting evidences: On the one hand, equilibrium studies of defect energy^{3,5,6,7}, Monte Carlo (MC) simulations of the root-mean-square current⁸, and resistance calculation⁹ have suggested that no finite- T ordering exists. The glass order parameter has furthermore been analytically shown to vanish at any finite T ¹⁰. On the other hand, the MC studies of the glass susceptibility¹¹, as well as dynamical simulations of the resistance^{12,13} and non-equilibrium relaxation¹⁴, have indicated a possibility of a finite- T transition. Also the MC simulations of the helicity modulus in Ref.¹⁵ were interpreted as being compatible with a finite- T transition.

In this paper we study a generalization of the 2D XY gauge glass model—the random gauge XY model—where both the temperature T and disorder strength r can be varied. When r has the maximum value 1, it corresponds to the usual XY gauge glass model, while in the opposite limit of $r = 0$, the standard XY model without disorder is recovered^{15,16,17}. It has been proposed that there is a Kosterlitz-Thouless (KT) like transition at a finite temperature T_c when the disorder strength is sufficiently small, and that as r is increased T_c becomes smaller until it vanishes as r reaches the critical disorder strength r_c (< 1) and $T_c = 0$ for $r > r_c$ ^{16,17}. However, even if the glass order parameter is zero and even if there is no finite- T KT transition for $r > r_c$, the existence of a finite- T transition with a different character cannot *a priori* be ruled out.

In the present paper we perform extensive MC simula-




FIG. 1: Sketch of the phase diagram in the disorder strength r and the temperature T (in units of J/k_B with the coupling strength J) plane. There exist two superconducting phases at low temperatures: the low- T KT phase marked by “SI” and the distinct phase “SII” for small r and large r , respectively. The high temperature normal phase is marked as “N”. The solid phase boundary represents the boundary of the low- T KT phase whereas the dashed represents the normal to superconducting transition for larger r .

tions to study the phase transition of the random gauge XY model. It is found that the system is superconducting at any r and that the transition from normal to superconducting phase is consistent with a Kosterlitz-Thouless (KT) type at weak disorder and a non-KT type at strong disorder. In addition to this we suggest that there exist two different superconducting phases at sufficiently low temperatures separated by a non-KT phase transition (see Fig. 1). For the special case of $r = 1$, which corresponds to the 2D XY gauge glass model, we also compute the root-mean-square current in the same way as in Ref.⁸ and, in contrast to Ref.⁸, again consistently find a finite-temperature transition.

II. MODEL AND SIMULATIONS

The Hamiltonian of the 2D random gauge XY model on an $L \times L$ square lattice under the fluctuating twist boundary condition (FTBC)¹⁹ is given by

$$\hat{H} = -J \sum_{\langle ij \rangle} \cos \left(\phi_{ij} \equiv \theta_i - \theta_j - \frac{1}{L} \mathbf{r}_{ij} \cdot \mathbf{\Delta} - A_{ij} \right), \quad (1)$$

where J is the coupling strength (set to unity from now on), the sum is over nearest neighbor pairs, $\mathbf{r}_{ij} \equiv \hat{\mathbf{x}}(\hat{\mathbf{y}})$ if $j = i + \hat{\mathbf{x}}(\hat{\mathbf{y}})$. The phase angle θ_i at the lattice point i satisfies the periodicity $\theta_{i+L\hat{\mathbf{x}}} = \theta_{i+L\hat{\mathbf{y}}} = \theta_i$, and $A_{ij} \in r[-\pi, \pi]$ is a uniform quenched random variable with the disorder strength $0 \leq r \leq 1$. The twist variable $\mathbf{\Delta} = (\Delta_x, \Delta_y)$ corresponds to the global twist across the system, i.e., the summation of the gauge invariant phase difference ϕ_{ij} along the x (y) direction equals Δ_x (Δ_y). For a given disorder realization, we first compute the distribution $P(\mathbf{\Delta})$, which is related to the free energy F by $\partial F / \partial \mathbf{\Delta} = -T(\partial \ln P / \partial \mathbf{\Delta})$. The twist variable $\mathbf{\Delta}_0$ which minimizes F (or maximizes P) is determined from P and then fixed when the helicity modulus $\Upsilon \equiv \partial^2 F / \partial \Delta^2$ and the 4th order modulus $\Upsilon_4 \equiv \partial^4 F / \partial \Delta^4$ are computed.

To ensure that the cooling is slow enough, we simultaneously cool two replicas (α and β) of the system and measure $\mathbf{\Delta}_0^{\alpha, \beta}$. For the first cooling at a new temperature we use 120000 update sweeps (for spin and twist variables respectively). Then we check that

$$|\Delta_{x,0}^{\alpha} - \Delta_{x,0}^{\beta}| < \delta \text{ and } |\Delta_{y,0}^{\alpha} - \Delta_{y,0}^{\beta}| < \delta \quad (2)$$

where δ sets the precision of the cooling. The idea with the annealing condition Eq. (2) is to keep the system close to the lowest-energy state at a particular temperature (so the system does not freeze into a local minimum that bias $[\Upsilon]$ and $[\Upsilon_4]$). Since the system moves more swiftly over the configuration space the higher the temperature is we can choose δ increasing with temperature. A choice that proves good in practice is

$$\delta = \begin{cases} 0.02\pi & \text{for } T < 0.3 \\ 0.15\pi & \text{for } 0.3 \leq T < 0.6 \\ \infty & \text{for } T \geq 0.6 \end{cases} \quad (3)$$

If the annealing condition Eq. (2) fail we repeat the cooling with three times as many update sweeps; if it fails again we increase the number of cooling sweeps a factor three again, and so on until the condition is fulfilled. When $\mathbf{\Delta}_0$ is chosen, before cooling, we let the system run until Eq. (2) is fulfilled with $\mathbf{\Delta}_0$ replaced by $\mathbf{\Delta}$.

We repeat the above calculations for more than 500 different disorder realizations (the disorder average is denoted by $[\dots]$ throughout this paper).

For the case of $r = 1$, we also use the periodic boundary condition PBC, corresponding to $\mathbf{\Delta} = 0$ in Eq. (1), and




FIG. 2: (a) The size dependence of the helicity modulus $[\Upsilon]$ at different temperatures T for the fully disordered case ($r = 1$)—the gauge glass model. At high T , $[\Upsilon]$ is shown to vanish as L is increased, whereas it saturates to a finite value at low T (qualitatively given by the dashed lines are linear extrapolations from the smallest sizes and correspond to power law behavior) (b) Finite-size scaling of $[\Upsilon]$ of the data in (a). From the standard finite-size scaling form, the critical temperature $T_c \approx 0.19$ and the critical exponent $\nu \approx 1.1$ are determined.

compute the root-mean-square current defined by⁸

$$I_{\text{rms}} \equiv \left[\left\langle \frac{1}{L} \sum_{\langle ij \rangle_x} \sin \phi_{ij} \right\rangle^2 \right]^{1/2}, \quad (4)$$

for comparisons with earlier works.

For I_{rms} , for each disorder, we use 50000 update sweeps for thermalizations and 500000 sweeps for measurements where the actual measurement was performed every tenth sweep. We used this computer time saving strategy since we found that performing the measurement every sweep gave closely the same result.




FIG. 3: The root-mean-square current I_{rms} . (a) The finite-size-scaling plot used in Ref.⁸ for $T_c = 0$ shows systematic deviations in the low-temperature region ($\nu = 2.2$ in Ref.⁸ is used). We include the data in Ref.⁸ for comparisons. (b) Finite-size scaling form in Eq. (6) yields $T_c \approx 0.2$ and $b \approx 0.5$. Inset: All data points in the main panel collapse to a smooth curve with $\nu = 1.1$, the same value as found for the helicity modulus in Fig. 2.

III. RESULTS FROM SIMULATIONS

We first investigate the standard 2D XY gauge glass model, corresponding to the fully disordered case ($r = 1$), and show in Fig. 2(a) the helicity modulus $[\Upsilon]$ as a function of the system size L for various temperatures. It is clearly shown that at high T , $[\Upsilon]$ goes to zero as the system size L is increased. The crucial point is that, at low enough temperatures ($T \lesssim 0.2$), $[\Upsilon]$ changes its curvature in terms of L , and appears to saturate to a finite value as L is increased. This behavior suggests a phase transition with a scale-invariant power law dependence at the critical temperature and a diverging characteristic length. Such a behavior can often be described by the finite-size scaling form

$$[\Upsilon] = L^{-b} f\left(L^{1/\nu}(T - T_c)\right), \quad (5)$$

where $[\Upsilon] \sim L^{-b}$ at the critical temperature T_c , and the critical exponent ν is related to the divergence of the coherence length. At T_c , the scaling function f has the same value irrespective of L , implying that $L^b[\Upsilon]$ versus T should have a unique crossing point at T_c for various sizes. In Fig. 2(b), it is clearly shown that $[\Upsilon]$ has a scale-invariant behavior $[\Upsilon] \propto L^{-0.4}$ at a unique T , signaling a phase transition. The inset of Fig. 2(b) furthermore confirms that this scaling behavior is consistent with the standard form (5) with $\nu \approx 1.1$.

Simple dimensional analysis for the non-disordered case, $r = 0$, gives for $[\Upsilon]$ the exponent $b = 0$ and from such a dimensional analysis one would likewise conclude that $b = 0$ also for the disordered case. Thus if $[\Upsilon]$ scales with $b \neq 0$ for the disordered case this is equivalent to the appearance of an anomalous dimension not accounted for by simple dimensional analysis. Our suggestion, based on the simulation results, is consequently that the disorder introduces such an anomalous dimension.

Figure 3 shows the root-mean-square current (4) for the PBC. We first note in Fig. 3(a) that the finite-size-scaling form for $T_c = 0$ used in Ref.⁸, $I_{\text{rms}} = L^{-1/\nu} f(T L^{1/\nu})$ with $\nu = 2.2$, shows systematic deviations from the data collapse to a single scaling curve at lower temperatures, in contrast to what was concluded in Ref.⁸, when more and better converged data are included. Furthermore a $T_c = 0$ -collapse cannot be achieved with any value of ν . On the other hand if we, in analogy with the finding for $[\Upsilon]$ above, use the scaling form

$$I_{\text{rms}} = L^{-b} f\left(L^{1/\nu}(T - T_c)\right), \quad (6)$$

which allows for the same anomalous dimension, one obtains the scaling plot in Fig. 3(b) with $b \approx 0.5$ and $T_c \approx 0.2$.¹⁸ From Figs. 2 and 3 we conclude that the 2D XY gauge glass ($r = 1$) exhibits a non-KT type superconducting to normal transition at $T_c \approx 0.2$ characterized by the existence of the anomalous dimension $b \approx 0.5$ and $\nu \approx 1.1$.

Next we investigate the phase transition with decreasing r . For $r = 0.9, 0.8, \dots, 0.5$, we obtain the finite-size scaling plots of the same quality as in Fig. 2(b), and determine the phase boundary (the dashed line in Fig. 1). As r is changed from 0.5 to 0.4 the exponents b and ν exhibit quite rapid changes, b from 0.27 to 0.06 and ν from 1.1 to 2.0. This, in our interpretation, reflects that near $r = 0.4$ the nature of the superconducting-normal transition changes from a non-KT type to a KT type.

For $r = 0$ the phase transition is of the KT nature and it has been suggested that this character should persist along the phase boundary up to some $r_c < 1$.^{16,17} The KT transition is characterized by that $[\Upsilon]$ jumps from a finite value $[\Upsilon] = 2T/\pi$ to zero as the phase line is crossed from the small (T, r) region. For $r = 0$ we find that the KT transition is also characterized by the increase of the 4th order modulus $|\Upsilon_4| \propto L^c$ at T_c with a positive exponent c , and $[\Upsilon_4]$ stays at a constant value below T_c as L is increased²⁰. Figure 4 (a) shows $[\Upsilon_4]$




FIG. 4: The fourth order modulus and helicity modulus (inset) as a functions of r for temperatures (a) 0.6 and (b) 0.1. The solid line in the inset represents the universal-jump condition for a KT transition $[\Upsilon] = 2T/\pi$.

for $T = 0.6$ as a function of r . The KT condition in the inset gives $r \approx 0.4$. This means that if the transition is of KT type then it would occur around $r \approx 0.4$. The onset of size dependence for $[\Upsilon_4]$ in Fig. 4 (a) is consistent with a transition at around $r \lesssim 0.4$. Thus the transition is compatible with a KT-character, although a different character cannot be ruled out. The situation for $T = 0.1$ in Fig. 4 (b) is quite different: the inset shows that the KT jump condition is not fulfilled and a KT transition can be ruled out. Yet there is a marked structure in $[\Upsilon]$ around $r \approx 0.4$. This structure corresponds to the onset of strong size dependence in $[\Upsilon_4]$. We interpret this onset as the reflection of a true divergence in $[\Upsilon_4]$ consistent with a phase transition. Thus we suggest that the whole boundary to the low- T KT phase (solid line in Fig. 1) is reflected in a divergence of $[\Upsilon_4]$ and that this line ends at $(T = 0, r_c \approx 0.4)$. A phase line which ends at $(T = 0, r_c \approx 0.4)$ has been found in many earlier investigations (see e.g. Ref.¹⁶ and references therein)²¹. The difference with earlier work is that in our case such a phase line is for lower temperatures between the two different superconducting phases SI and SII (as demonstrated by our direct calculation of $[\Upsilon]$), whereas earlier work have

concluded that such a phases line does indeed exist but separates a superconducting phase from a normal phase all the way down to $T = 0$. Our suggested phase diagram is thus consistent with earlier work as to the existence of a phase line ending at $(T = 0, r_c \approx 0.4)$. For temperatures larger than the merging with the second phase line (above dashed line in Fig. 1) the transition is consistent with a KT transition although a different character cannot be ruled out. However, below the merging with this second line the transition is not a KT transition. It may be that there still is a jump in $[\Upsilon]$ at this transition, but this is then between two non-vanishing values.

Based on the numerical evidences we suggest the structure of the phase diagram is sketched in Fig. 1. One striking feature is the finite- T transition line between normal and superconducting phases, starting from the boundary of the low- T KT-phase (solid line) and changing into the (dashed) line which ends at $T_c \approx 0.2$ at $r = 1$. The latter line (dashed line) is characterized by the appearance of an anomalous dimension b . Since the KT transition does not have such an anomalous dimension it follows that, if the transition along the boundary to the low- T KT phase boundary has KT character, then b should approach $b = 0$ when the two transition line merge. Thus the rapid drop from $b \approx 0.27$ to $b \approx 0.06$ as r is changed from $r = 0.5$ to 0.4 is consistent with a change over to a KT transition. Another interesting feature is that the phase line (solid line in Fig. 1), associated with the divergence of $[\Upsilon_4]$, continues even inside the superconducting region and ends at $r \approx 0.4$ for $T = 0$. Although the transition separating the KT phase and the normal phase (SI and N, respectively, in Fig. 1), may well be of the true KT type, manifested by a universal jump in the helicity modulus, the phase line separating SI and SII in Fig. 1 is not a KT transition and the helicity modulus has a non-zero value on both sides.

It is interesting to note that earlier works have found evidences for only two phases separated by a *single* phase line; either, in the more prevailing view, a phase line ending at a point $T = 0$ for a finite $r_c < 1$, or, in the less prevailing view, a phase line ending at a point $T > 0$ for $r = 1$. From our numerical simulations we instead suggest three distinct phases separated by *two* phase lines, which combines the two earlier proposed scenarios and provides a unified picture: On the one hand, in Ref.⁵, the end point of the phase boundary to the low- T KT-phase has been obtained to be $r_c \approx 0.37$ and $T = 0$, which is consistent with our cruder estimate $r \approx 0.4$. On the other hand, the phase transition point at $T \approx 0.22$ with $\nu = 1.1$, found in Refs.¹¹ and¹³ for $r = 1$, is in very good agreement with the end point of our finite- T phase line ($T \approx 0.2, \nu \approx 1.1$). However, the difference with the previous work⁵ is that according to our interpretation the phase line below $T \approx 0.2$ ending at $T = 0$ and $r_c \approx 0.4$ separates two distinct superconducting phases, while the whole phase line in Ref.⁵ is for the superconducting to normal transition. A very hand-waving picture of the scenario of our phase diagram in terms of vortex motion

is sketched in Fig. 1: In SI the vortex motion is suppressed by vortex pair binding, in N pair unbinding has occurred and free vortices exists which are not entirely pinned by the disorder, whereas in SII vortex pair unbinding has occurred but the vortices are pinned by the disorder.

IV. SUMMARY

One main conclusion from this phase diagram is that the XY gauge glass model ($r = 1$) has a finite- T transition. How is this possible in view of earlier conflicting evidences? In Ref.⁸ it was concluded, on the basis of an analysis of data for the root-mean-square current for standard periodic boundary conditions²², that no finite- T transition exists in 2D. We have found that, taking into account the possibility of an anomalous scaling dimension, I_{rms} displays a transition at a finite-temperature (see Fig. 3)²⁰. Another puzzling evidence to the contrary

is the $T = 0$ calculations of the size scaling of the defect energy^{3,5,6,7}. However, as discussed in Refs.⁷ and²³, the local vorticity conservation must be properly taken into account when calculating the energy barriers. Thus the energy barrier for vortex dissipation increases with system size when taking the local vorticity conservation into account²³. This growing of the energy barrier for vortex dissipation with system size supports the possibility of a finite- T transition²³.

The appearance of a new superconducting phase for the XY random gauge model is intriguing. In particular, since it is neither a low T KT phase nor a phase with a finite glass order parameter. The true nature of this phase and the existence of similar phases in other related models are open questions.

Support from the Swedish Natural Research Council through Contract No. F 5102-659/2001 is gratefully acknowledged. B.J.K. was supported by the Korea Science and Engineering Foundation through Grant No. R14-2002-062-01000-0.

-
- * Electronic address: holme@tp.umu.se
 - † Electronic address: beomjun@ajou.ac.kr
 - ‡ Electronic address: minnhagen@nordita.dk
 - ¹ D. A. Huse and H. S. Seung, Phys. Rev. B **42**, 1059 (1990).
 - ² M. P. A. Fisher, Phys. Rev. Lett. **62**, 1415 (1989); D. S. Fisher, M. P. A. Fisher, and D. A. Huse, Phys. Rev. B **43**, 130 (1991).
 - ³ M. Cieplak, J. R. Banavar, and A. Khurana, J. Phys. A **3**, L145 (1991); M. Cieplak *et al.*, Phys. Rev. B **45**, 786 (1992).
 - ⁴ J. D. Reger *et al.*, Phys. Rev. B **44**, 7147 (1991); C. Wengel and A. P. Young, Phys. Rev. B **56**, 5918 (1997).
 - ⁵ J. M. Kosterlitz and M. V. Simkin, Phys. Rev. Lett. **79**, 1098 (1997).
 - ⁶ M. J. P. Gingras, Phys. Rev. B **45**, 7547 (1992); H. S. Bokil and A. P. Young, Phys. Rev. Lett. **74**, 3021 (1998); J. M. Kosterlitz and M. Akino, Phys. Rev. Lett. **81**, 4672 (1998).
 - ⁷ M. P. A. Fisher, T. A. Tokuyasu, and A. P. Young, Phys. Rev. Lett. **66**, 2931 (1991).
 - ⁸ J. D. Reger and A. P. Young, J. Phys. A **26**, L1067 (1993).
 - ⁹ R. A. Hyman *et al.*, Phys. Rev. B **51**, 15304 (1995).
 - ¹⁰ H. Nishimori, Physica A **205**, 1 (1994).
 - ¹¹ M. Y. Choi and S. Y. Park, Phys. Rev. B **60**, 4070 (1999).
 - ¹² Y. H. Li, Phys. Rev. Lett. **69**, 1819 (1992).
 - ¹³ B. J. Kim, Phys. Rev. B **62**, 644 (2000).
 - ¹⁴ B. J. Kim *et al.*, Phys. Rev. B **56**, 6007 (1997).
 - ¹⁵ G. S. Jeon, S. Kim, and M. Y. Choi, Phys. Rev. B **51**, 16211 (1995).
 - ¹⁶ J. Maucourt and D. R. Grempel, Phys. Rev. B **56**, 2572 (1997).
 - ¹⁷ M. Rubinstein, B. Shraiman, and D. R. Nelson, Phys. Rev. B **27** 1800 (1983); T. Natterman *et al.*, J. Phys. (France) I **5**, 565 (1995); M.-C. Cha and H. A. Fertig, Phys. Rev. Lett. **74**, 4867 (1995); L.-H. Tang, Phys. Rev. B **54**, 3350 (1996).
 - ¹⁸ Preliminary results for parallel tempering Monte Carlo give the same qualitative result.²⁰
 - ¹⁹ P. Olsson, Phys. Rev. B **46**, 14598 (1992); **52**, 4526 (1995).
 - ²⁰ For more details, B. J. Kim, P. Minnhagen, and P. Holme (unpublished).
 - ²¹ This boundary of SI is completely in accord with the change from power law to exponential decay of the spin correlation function found in Ref.16²⁰. In SII the higher order correlation function discussed in Ref.15 has a power-law decay²⁰.
 - ²² With the FTBC, the r.m.s. current is identically zero.
 - ²³ P. Holme and P. Olsson, Europhys. Lett. **60**, 439 (2002).